



# Study of Fuzzy system to Various Objective Decision Making Models

Anil Kumar<sup>1</sup>, Priyadarshan<sup>2</sup>

Assistant Professor, Ch. Devilal University, Sirsa, Haryana <sup>1,2</sup>  
Email: anilchawla373@gmail.com, pdbeniwal40@gmail.com

Abstract- Ranking of Intuitionist Fuzzy Sets (IFSs) and Interval Valued Intuitionist Fuzzy Sets (IVIFSs) are very often required in decision making. Various decision making models are already in literature. Shiny Jose and Sunny Kuriakose in 2018 introduced a score function for ranking interval valued intuitionist fuzzy numbers. In this paper I have proposed a decision making model in multi person decision making problems in IVIF environment.

**Keywords:** Fuzzy, Sets, decision, models

## 1. INTRODUCTION

Following the introduction of Fuzzy set (FS for short) by L. A. Zadeh in 1965, Krassimir Atanassov introduced the notion of IFS which has been found a better tool to model decision problems [1]. Multicriteria decision making models based on IFS theoretical tools were introduced in the decision theory by Z. S. Xu [2,3]. This was extended to IVIFS [4]. Later many researchers studied the problem of ranking IFSs. Shiny Jose and Sunny Kuriakose introduced a score function in [5] to rank the alternatives in IVIF context. In this paper I consider the situation of multiperson multicriteria decision making in IVIF context, and solve the problem using the accuracy function given in [5]. Section 2 contains basic definitions and results. Section 3 contains the required score function. New decision making model and its illustration are given in section 4.

## 2. PRELIMINARIES

**Definition 2.1. [1] Intuitionistic Fuzzy Sets.** Let  $X$  be a given set. An Intuitionistic fuzzy set  $A$  in  $X$  is given by,

$$A = \{(x, \mu_A(x), \nu_A(x)) / x \in X\}$$

Where

$\mu_A, \nu_A : X \rightarrow [0, 1]$ , and  $0 \leq \mu_A(x) + \nu_A(x) \leq 1$ ,  $\mu_A(x)$  is the degree of membership of the element  $x$  in  $A$  and  $\nu_A(x)$  is the degree of non-membership of  $x$  in  $A$ . For each  $x \in X$ ,  $\pi_A(x) = 1 - \mu_A(x) - \nu_A(x)$  is called the degree of hesitation.

**Definition 2.2. [3] Interval valued intuitionistic fuzzy sets.** Let  $D [0, 1]$  be the set of all closed subintervals of the interval  $[0, 1]$ . Let  $X \neq \emptyset$  be a given set.

An interval valued intuitionistic fuzzy set  $A$  in  $X$  is given by

$$A = \{(x, \mu_A(x), \nu_A(x)) : x \in X\},$$

where  $\mu_A: X \rightarrow D [0, 1]$ ,  $\nu_A: X \rightarrow D [0, 1]$  with the condition

$$0 \leq \sup_x \mu_A(x) + \sup_x \nu_A(x) \leq 1$$

The intervals  $\mu_A(x)$  and  $\nu_A(x)$  denote, respectively, the degree of belongingness and the degree of non-belongingness of the element  $x$  to the set  $A$ . Thus, for each  $x \in X$ ,  $\mu_A(x)$  and  $\nu_A(x)$  are closed intervals whose lower and upper end points are respectively, denoted by  $\mu_{AL}(x)$ ,  $\mu_{AU}(x)$  and  $\nu_{AL}(x)$ ,  $\nu_{AU}(x)$ .

$A$  can also be denoted by,

$A = \{(x, [\mu_{AL}(x), \mu_{AU}(x)], [\nu_{AL}(x), \nu_{AU}(x)]) : x \in X\}$ ,  
Where  $0 \leq \mu_{AU}(x) + \nu_{AU}(x) \leq 1$ ,  $\mu_{AL}(x) \geq 0$  and  $\nu_{AL}(x) \geq 0$ . For every element  $x \in X$ , the hesitancy degree of an intuitionistic fuzzy interval of  $x \in X$  in  $A$  is defined as

$$\pi_A(x) = 1 - \mu_A(x) - \nu_A(x) = [1 - \mu_{AU}(x) - \nu_{AU}(x), 1 - \mu_{AL}(x) - \nu_{AL}(x)].$$

We will denote the set of all the IVIFS in  $X$  by  $IVIFS(X)$ .

**Definition 2.3. [4] Weighted geometric average operator for IVIFSs.** Let  $A_j (j = 1, 2, \dots, n) \in IVIFS(X)$ . The weighted geometric average operator is defined by

$$G_w(A_1, A_2, A_3, \dots, A_n) = \Pi A_j^{w_j} = ([\Pi \mu_{Aj}^{Lw_j}(x), \Pi \mu_{Aj}^{Uw_j}(x)], [1 - \Pi (1 - \nu_{AjL}(x))^{w_j}, 1 - \Pi (1 - \nu_{AjU}(x))^{w_j}])$$

Where  $w_j$  is the weight of  $A_j (j = 1, 2, \dots, n)$ ,  $w_j \in [0, 1]$  and  $\sum^n w_j = 1$ .

Assume  $w = 1/n (j = 1, 2, \dots, n)$  then  $G$  for  $A_1, A_2, \dots, A_n$ . Clearly  $G_w$  is an IVIFS.

**Definition 2.4. [4] Weighted arithmetic average operator for IVIFSs.** Let  $A_j (j = 1, 2, \dots, n) \in IVIFS(X)$ . The weighted arithmetic average operator is defined by:



$$F_w(A_1, A_2, A_3 \dots A_n) = \sum_{j=1}^n w_j A_j = ([1 - \prod_{j=1}^n (1 - \mu_{AjL}(x))^{w_j}], [1 - \prod_{j=1}^n (1 - \mu_{AjU}(x))^{w_j}], [\prod_{j=1}^n v_{AjL}^{w_j}(x), \prod_{j=1}^n v_{AjU}^{w_j}(x)])$$

**3 SCORE FUNCTION**

For an interval valued intuitionist fuzzy number  $A_1$  obtained by the weighted geometric average operator, say  $A_1(a_1, b_1, c_1, d_1)$ , the score function  $S$  at  $A_1$  based on the hesitancy degree denoted by  $S(A_1)$  is defined by,

$$S(A_1) = \frac{(a_1 + b_1 - c_1 d_1)}{2}$$

**Result 3.1. [5]** For any interval-valued intuitionist fuzzy subset  $A = ([a, b], [c, d])$ , the new proposed score  $S(A) \in [-1, 1]$ .

**Result 3.2. [5]** For any two comparable interval valued intuitionist fuzzy sets  $A$  and  $B$ , if  $A \subseteq B$  then  $S(A) \leq S(B)$ .

**Result 3.3. [5]** If  $A = [a_1, b_1], [c_1, d_1]$  and  $B = [a_2, b_2], [c_2, d_2]$  be two interval valued intuitionist fuzzy sets such that  $S(A) = S(B)$ , then  $A = B$ .

**4 MULTI-PERSON DECISION MAKING MODEL IN IV IF ENVIRONMENT**

Xu in [6] elicit a method for group decision making, where any piece of information provided by the decision makers is expressed as intuitionist fuzzy decision matrices and the information about attribute weights is partially known, or may be constructed by various methods. They first use the intuitionist fuzzy hybrid geometric (IFHG) operator to aggregate all individual intuitionist fuzzy decision matrix provided by the decision makers into the collective intuitionist fuzzy decision matrix. Then they utilize the score function to calculate the score value of each attribute and construct the score matrix of the collective intuitionist fuzzy decision matrix. Based on the score matrix and the given information about attribute weight, they establish an optimization model to determine the weights of attributes. After that they use the obtained attribute weights and the intuitionist fuzzy weighted geometric (IFWG) operator to aggregate the intuitionist fuzzy information in the collective intuitionist fuzzy decision matrix to get

overall intuitionist fuzzy values of alternatives by which the ranking of all the given alternatives can be obtained.

I apply this method to interval valued intuitionist fuzzy decision making model, but in a different way as follows.

**Computational procedure**

**Step 1.**

First we calculate intuitionist fuzzy weighted geometric average value for each alternatives using definition 2.3 Here we assume that criteria weights are known.

**Step 2.**

Find the score for each alternatives, by the score function in section 3, i.e., Obtain the collective score matrix as follows.

	$A_1$	$A_2$	...	$A_m$
$D_1$	$S_{11}$	$S_{12}$	...	$S_{1m}$
$D_2$	$S_{21}$	$S_{22}$	...	$S_{2m}$
.	.	.	.	.
$D_n$	$S_{n1}$	$S_{n2}$	...	$S_{nm}$

**Step 3.** Let  $W_{Di}$  be the weight for the decision maker  $D_i, i = 1, 2, \dots, n$ . Then find the collective score for each alternative  $A_j (j = 1, 2, \dots, m)$  by the formula

$$S(A_j) = \sum_{i=1}^n W_{Di} S_{ij}$$

**Step 4.** Rank the alternatives according to their collective score. Now we illustrate this with an example.

**Illustrative example**

In a youth festival, three decision makers have to rank four dancers ( $A_1, A_2, A_3, A_4$ ) based on five criteria ( $C_1, C_2, C_3, C_4, C_5$ ). The decision of the decision makers are given respectively in the following decision matrices.

	$A_1$	$A_2$	$A_3$	$A_4$
$C_1$	[.2, .3], [.4, .5]	[.7, .8], [.1, .15]	[.6, .7], [.1, .2]	[.2, .3], [.4, .5]
$C_2$	[.6, .7], [.2, .3]	[.5, .6], [.2, .3]	[.6, .7], [.1, .3]	[.3, .4], [.5, .6]
$C_3$	[.5, .6], [.2, .3]	[.4, .5], [.3, .4]	[.5, .6], [.1, .2]	[.5, .6], [.1, .2]
$C_4$	[.4, .6], [.2, .3]	[.6, .7], [.1, .2]	[.6, .7], [.1, .2]	[.6, .6], [.1, .2]
$C_5$	[.5, .6], [.1, .2]	[.4, .5], [.1, .2]	[.5, .6], [.1, .2]	[.4, .5], [.2, .3]

**Decision Matrix -1**

	A <sub>1</sub>	A <sub>2</sub>	A <sub>3</sub>	A <sub>4</sub>
C <sub>1</sub>	[.1, .3], [.4, .5]	[.6, .7], [.1, .2]	[.7, .8], [.1, .2]	[.2, .3], [.5, .6]
C <sub>2</sub>	[.6, .7], [.1, .2]	[.7, .8], [.2, .3]	[.6, .7], [.1, .2]	[.3, .4], [.5, .6]
C <sub>3</sub>	[.4, .5], [.2, .3]	[.4, .5], [.3, .4]	[.6, .7], [.1, .3]	[.5, .6], [.1, .2]
C <sub>4</sub>	[.4, .6], [.2, .3]	[.5, .6], [.2, .3]	[.5, .6], [.1, .2]	[.3, .4], [.1, .2]
C <sub>5</sub>	[.4, .6], [.1, .2]	[.4, .5], [.1, .2]	[.4, .5], [.1, .2]	[.4, .5], [.1, .2]

Decision Matrix -2

	A <sub>1</sub>	A <sub>2</sub>	A <sub>3</sub>	A <sub>4</sub>
C <sub>1</sub>	[.2, .3], [.4, .5]	[.6, .7], [.1, .2]	[.3, .4], [.4, .5]	[.1, .2], [.4, .5]
C <sub>2</sub>	[.7, .8], [.1, .2]	[.5, .6], [.2, .3]	[.3, .4], [.4, .5]	[.3, .4], [.5, .6]
C <sub>3</sub>	[.4, .5], [.2, .3]	[.4, .5], [.3, .4]	[.6, .7], [.1, .2]	[.4, .5], [.1, .2]
C <sub>4</sub>	[.4, .6], [.2, .3]	[.5, .7], [.1, .2]	[.3, .4], [.1, .2]	[.6, .7], [.1, .2]
C <sub>5</sub>	[.4, .6], [.1, .2]	[.3, .4], [.4, .5]	[.4, .5], [.2, .3]	[.4, .5], [.2, .3]

Decision Matrix -3

Assuming weights for criteria C<sub>1</sub>, C<sub>2</sub>, C<sub>3</sub>, C<sub>4</sub> and C<sub>5</sub> as 0.3, 0.2, 0.15, 0.2 and 0.15 respectively, we

obtain weighted geometric average value for A<sub>j</sub> as follows.

#### Aggregated Matrix

	A <sub>1</sub>	A <sub>2</sub>	A <sub>3</sub>	A <sub>4</sub>
D <sub>1</sub>	[.38, .50], [.25, .35]	[.54, .64], [.15, .24]	[.57, .67], [.1, .23]	[.34, .44], [.31, .41]
D <sub>2</sub>	[.29, .49], [.24, .34]	[.53, .63], [.17, .28]	[.57, .67], [.1, .22]	[.29, .40], [.33, .43]
D <sub>3</sub>	[.36, .50], [.24, .34]	[.47, .59], [.20, .30]	[.35, .45], [.28, .38]	[.27, .39], [.30, .41]

We know that each weighted geometric average value is an IVIFN. So we can find the collective

score for each alternative as given in step-3. It is elicited in the table given below.

#### Collective Score Matrix

	A <sub>1</sub>	A <sub>2</sub>	A <sub>3</sub>	A <sub>4</sub>
D <sub>1</sub>	.396	.572	.609	.326
D <sub>2</sub>	.349	.556	.609	.274
D <sub>3</sub>	.389	.5	.347	.268

Let the weight for each decision maker be as follows. Weight for decision maker-1, W<sub>D1</sub> = 0.4

Weight for decision maker-2, W<sub>D2</sub> = 0.3

Weight for decision maker-3, W<sub>D3</sub> = 0.3

Thus the collective score for each alternative be,  
S(A<sub>1</sub>) = (0.4 × 0.396) + (0.3 × 0.349) + (0.3 × 0.389),

Similarly we can find the score for other alternatives also.

i.e., S(A<sub>1</sub>)=0.3798, S(A<sub>2</sub>) =0.5456, (A<sub>3</sub>) =0.5304 and S(A<sub>4</sub>)=0.293.

Thus we rank the alternatives according to their score as follows,

A<sub>2</sub>> A<sub>3</sub>> A<sub>1</sub>> A<sub>4</sub>

#### 5 CONCLUSION

In this paper I have introduced a decision making model which is elegant than the existing models in literature. This can be used in Multi person multi criteria decision making problems in interval valued intuitionistic fuzzy environment.

#### REFERENCES

- [1] Atanassov, K. T., 1986, "Intuitionistic fuzzy sets," Fuzzy Sets and Systems, 20(1), pp. 87–96.
- [2] Xu, Z. S., Yager, R. R., 2006, "Some geometric aggregation operators based on intuitionistic fuzzy environment," International Journal of General Systems, Vol. 33, pp. 417–433.
- [3] Xu, Z. S., 2007, "Intuitionistic fuzzy



*International Journal of Engineering, Pure and Applied Sciences,*  
*Vol. 4, No. 1, March-2019*

- aggregation operators,” IEEE Transactions on Fuzzy Systems, Vol. 15, pp. 1179–1187.
- [4] Xu, Z. S., 2007, “Methods for aggregating interval valued intuitionistic fuzzy information and their application to decision making”, Control and Decision, 22(2), pp. 215–219.
- [5] Shiny Jose, Sunny Kuriakose, 2013, Multi criteria Decision Making under Interval Valued Intuitionistic Fuzzy Context, The Journal of Fuzzy Mathematics, Vol. 21, No. 4, pp. 37–41.
- [6] Xu, Z.S., “Multiperson multiattribute decision making models under intuitionistic fuzzy environment”, Fuzzy Optimization and Decision Making, Vol. 6, 2007, pp. 221–236