



EVALUATION OF ANTIOXIDANT, ANTIMICROBIAL, AND ANTI-INFLAMMATORY ACTIVITIES OF SELECTED TRADITIONALLY USED MEDICINAL PLANTS

Jatinder Singh¹ & Dr. Jitendra Gupta²

Research Scholar, Arni University
 Professor, Arni University, Indora , H.P

ABSTRACT

Traditionally used medicinal plants constitute an important source of bioactive compounds with potential antioxidant, antimicrobial and anti-inflammatory properties. Their therapeutic relevance is frequently associated with secondary metabolites, including phenolic acids, flavonoids, tannins, alkaloids, terpenoids and saponins. The present research paper proposes a comparative evaluation of five medicinal plants, namely *Withania somnifera* (L.) Dunal, *Tinospora cordifolia* (Willd.) Hook. f. & Thomson, *Ocimum tenuiflorum* L. (syn. *Ocimum sanctum* L.), *Asparagus racemosus* Willd. and *Terminalia chebula* Retz. The study framework integrates pharmacognostical authentication, standardised extraction, preliminary phytochemical screening, quantitative estimation of total phenolic and flavonoid contents, and in vitro assessment of antioxidant, antimicrobial and anti-inflammatory activities. Antioxidant activity may be investigated using DPPH and ABTS radical-scavenging assays, antimicrobial efficacy through agar diffusion and minimum inhibitory concentration methods, and anti-inflammatory potential through protein-denaturation or other validated experimental models. Comparative tables and illustrative graphical datasets are included to demonstrate how experimental findings can be organised and interpreted. These numerical examples are hypothetical and are not presented as observed laboratory results. The proposed framework provides a basis for identifying promising plant extracts, examining relationships between phytochemical composition and biological activity, and establishing quality-control parameters for further pharmacological investigation. Rigorous experimental validation, toxicity assessment and chemical characterisation remain essential before therapeutic applications can be recommended.

Keywords: Medicinal plants; antioxidant activity; antimicrobial activity; anti-inflammatory activity; phytochemical screening; DPPH assay; minimum inhibitory concentration; traditional healthcare; herbal drug standardisation.

INTRODUCTION

Medicinal plants have occupied a central position in traditional healthcare systems, including Ayurveda and other indigenous medical traditions, where plant-derived preparations are used to manage a wide range of health conditions. Their continued use has encouraged scientific investigation into the chemical constituents responsible for their biological effects. Modern pharmacognosy and experimental pharmacology provide methods for authenticating plant materials, characterising their phytochemical profiles and evaluating biological activities under controlled conditions.

Oxidative stress occurs when the production of reactive oxygen and nitrogen species exceeds the capacity of endogenous antioxidant defence systems. Excessive oxidative stress can damage cellular lipids, proteins and nucleic acids and is implicated in several pathological processes. Plant-derived phenolics and flavonoids can exhibit antioxidant activity through mechanisms such as radical scavenging, metal chelation and modulation of cellular defence pathways. Nevertheless, antioxidant activity observed in a chemical assay does not necessarily establish efficacy

within living organisms, where absorption, metabolism and tissue distribution influence biological outcomes (Halliwell & Gutteridge, 2015).

Antimicrobial resistance has also increased interest in natural products as sources of compounds that may inhibit pathogenic microorganisms. Medicinal plants contain chemically diverse constituents capable of affecting microbial membranes, enzymes, cellular metabolism and biofilm formation. However, antimicrobial effects depend on the plant species, extraction solvent, concentration, test organism and experimental conditions. In vitro inhibition alone does not demonstrate clinical effectiveness or establish that a plant preparation can replace conventional antimicrobial treatment.

Inflammation is a protective physiological response to infection, tissue injury and other harmful stimuli. When inadequately regulated or persistent, inflammatory processes can contribute to tissue damage and chronic disease. Several plant-derived constituents have demonstrated anti-inflammatory effects in experimental systems through mechanisms associated with inflammatory mediators, oxidative



pathways and cellular signalling. Yet these mechanisms require careful validation, particularly when results from isolated biochemical systems are extrapolated to complex biological processes.

The present study focuses on five traditionally used medicinal plants: *Withania somnifera*, *Tinospora cordifolia*, *Ocimum tenuiflorum*, *Asparagus racemosus* and *Terminalia chebula*. These species represent different botanical groups and phytochemical profiles, making them suitable candidates for comparative investigation. Their selection also permits examination of how differences in plant material and extraction characteristics may relate to measured biological responses.

A systematic comparative framework is necessary because results obtained from different studies are often difficult to compare owing to variation in extraction procedures, assay protocols, reference standards and reporting units. Standardised experimental procedures, authenticated botanical material, appropriate controls and transparent statistical analysis can improve reproducibility and support more meaningful interpretation.

MATERIALS AND METHODS

Study design and selection of medicinal plants

A comparative, laboratory-based experimental design is proposed to evaluate the three biological activities of the selected plant extracts. The investigation should use authenticated plant material, standardised extraction procedures and replicated assays to reduce experimental variability. The selected species, plant parts and principal research rationale are presented in Table 1.

Table 1. Selected medicinal plants and their relevance to the proposed investigation

Scientific name	Plant part proposed	Major phytochemical groups of interest	Research focus
<i>Withania somnifera</i>	Roots	Withanolides, alkaloids, phenolic compounds	Antioxidant and anti-inflammatory potential
<i>Tinospora cordifolia</i>	Stems	Diterpenoids, alkaloids, glycosides, phenolics	Antioxidant and immune-related activity
<i>Ocimum tenuiflorum</i>	Leaves	Phenolics, flavonoids, volatile constituents	Antioxidant and antimicrobial potential
<i>Asparagus racemosus</i>	Roots	Steroidal saponins, polysaccharides, phenolics	Antioxidant and anti-inflammatory potential
<i>Terminalia chebula</i>	Fruits	Hydrolysable tannins, phenolic acids, flavonoids	Antioxidant and antimicrobial potential

Collection, authentication and preparation

Plant materials should be obtained from documented sources and authenticated by a qualified botanist or pharmacognosist. Voucher specimens should be deposited in an appropriate herbarium, with accession numbers recorded in the final manuscript. Collection location, season, plant maturity and plant-part identity should be documented because these variables can influence phytochemical composition.

The samples should be cleaned, dried under controlled conditions, pulverised and stored in airtight containers protected from moisture and light. Macroscopic and microscopic characteristics, foreign matter, moisture content, total ash, acid-insoluble ash and extractive values should be examined where applicable. These parameters help assess botanical identity and material quality, following recognised medicinal-plant quality-

control guidance (World Health Organization [WHO], 1998).

Extraction and phytochemical screening

A consistent extraction procedure should be used for all five plants. A suitable starting protocol involves extracting a measured quantity of powdered plant material with a defined volume of ethanol–water or another justified solvent system. The solvent concentration, extraction duration, temperature, extraction cycles and solvent-to-material ratio must remain consistent or be explicitly documented.

After filtration and solvent removal under controlled conditions, extraction yield should be calculated as:

$$\text{Extraction yield (\%)} = \frac{\text{Mass of dried extract}}{\text{Mass of dry plant material}} \times 100$$

Preliminary phytochemical screening may examine phenolics, flavonoids, tannins, alkaloids, saponins, terpenoids and glycosides using appropriate validated procedures. Total phenolic content may be estimated by the Folin–Ciocalteu method and expressed as milligrams of gallic acid equivalents per gram of dried extract (mg GAE/g). Total flavonoid content may be determined using a suitable validated colorimetric method and expressed as milligrams of quercetin equivalents per gram (mg QE/g). Appropriate blanks and calibration curves are required.

Where facilities permit, HPTLC, HPLC or LC–MS can be used to characterise selected constituents. Preliminary colour reactions alone should not be considered definitive evidence of chemical identity.

Evaluation of antioxidant activity

The DPPH radical-scavenging assay can be used to determine the capacity of extracts to reduce a stable free radical. The method is based on a change in absorbance following reaction with antioxidant constituents (Blois, 1958).

Extracts should be tested over a concentration range, with a suitable positive control such as ascorbic acid or Trolox. After correction for extract colour and reagent background, percentage radical-scavenging activity can be calculated as:

$$\text{Scavenging (\%)} = \frac{A_c - A_s}{A_c} \times 100$$

The concentration producing 50% radical-scavenging activity (IC₅₀) should be estimated from an appropriate concentration–response model. A lower IC₅₀ indicates stronger activity under the specified assay conditions. ABTS radical-scavenging assays can provide complementary evidence, although results from different assays should not be treated as interchangeable.

Evaluation of antimicrobial activity

Antimicrobial screening should employ appropriately characterised reference strains and laboratory-



approved biosafety procedures. Selected Gram-positive and Gram-negative bacteria may be investigated, subject to institutional approval and laboratory capability. Yeast or fungal testing can be added if a separately validated protocol is available.

Agar diffusion can provide preliminary information through measurement of inhibition-zone diameters. However, zone size depends on diffusion, solubility and other factors; therefore, it cannot independently establish comparative potency. Broth microdilution should be used to determine the minimum inhibitory concentration (MIC), defined as the lowest tested concentration preventing visible growth under the specified conditions.

The inoculum, medium, incubation conditions, solvent controls, growth controls and reference antimicrobial agents should be standardised. CLSI M07 provides established procedures for dilution-based antimicrobial susceptibility testing of aerobic bacteria, although plant-extract testing may require additional validation because crude mixtures can interfere with conventional endpoints.

Evaluation of anti-inflammatory activity

The proposed screening strategy may include inhibition of heat-induced protein denaturation as an initial in vitro model. The assay measures the ability of an extract to reduce a defined change in protein structure under controlled experimental conditions. Percentage inhibition may be calculated using corrected control and sample absorbance values:

$$\text{Inhibition (\%)} = \frac{A_c - A_s}{A_c} \times 100$$

The model requires suitable vehicle controls, a positive reference compound and correction for extract colour or turbidity. It should be complemented, where feasible, by more biologically informative assays, such as measurement of inflammatory mediators in validated cell-based models.

Protein-denaturation inhibition is a screening endpoint rather than direct evidence of anti-inflammatory efficacy in humans. Further investigation should address cytotoxicity, concentration-dependent responses and potential interference with the assay.

Statistical analysis

Each assay should include independent experimental replicates, with the final number determined by an appropriate study design and power analysis. Technical replicates should not be mistaken for independent biological samples. Results may be summarised as mean $\pm$ standard deviation, with concentration-response modelling used to estimate IC₅₀ values.

One-way analysis of variance followed by a suitable multiple-comparison procedure may be used when assumptions are satisfied. Otherwise, appropriate non-parametric methods should be considered. Correlation analysis can examine relationships between phenolic

content, flavonoid content and biological endpoints. Statistical significance should be interpreted alongside effect sizes, confidence intervals and experimental limitations.

RESULTS AND ILLUSTRATIVE DATA PRESENTATION

Data-status statement: No laboratory dataset was supplied with the request. Accordingly, the following numerical results are explicitly hypothetical and are included to demonstrate research-paper presentation, tables and graph construction. They must not be represented as experimental observations, validated findings or published measurements. Actual results should replace these examples before the manuscript is submitted as an original experimental research paper.

Comparative phytochemical and antioxidant profile

Table 2 presents an illustrative dataset for total phenolic content, total flavonoid content and DPPH radical-scavenging activity.

Medicinal plant	Total phenolics (mg GAE/g)	Total flavonoids (mg QE/g)	DPPH IC ₅₀ (µg/mL)
<i>W. somnifera</i>	68.4	31.2	74.5
<i>T. cordifolia</i>	82.6	37.8	61.3
<i>O. tenuiflorum</i>	96.8	45.1	42.7
<i>A. racemosus</i>	54.2	24.6	89.1
<i>T. chebula</i>	128.5	52.4	28.6

In this illustrative dataset, *T. chebula* has the highest estimated total phenolic and flavonoid contents and the lowest DPPH IC₅₀. *O. tenuiflorum* ranks second on these illustrative measures. This pattern would be consistent with a possible association between phenolic abundance and radical-scavenging activity, but it would not establish a causal relationship. Correlation analysis using actual independent samples would be required to assess the association, and additional chemical or biological mechanisms could contribute to the observed response.

Comparative antimicrobial activity

Antimicrobial activity should be assessed against selected reference microorganisms using both a preliminary diffusion assay and a quantitative dilution method. Table 3 provides an example of how inhibition-zone measurements could be reported for a selected bacterial strain.

Plant extract	Test organism	Inhibition zone (mm)	Illustrative MIC (µg/mL)
<i>W. somnifera</i>	<i>Staphylococcus aureus</i>	12.4	500
<i>T. cordifolia</i>	<i>S. aureus</i>	14.1	250
<i>O. tenuiflorum</i>	<i>S. aureus</i>	18.6	125
<i>A. racemosus</i>	<i>S. aureus</i>	10.8	1,000
<i>T. chebula</i>	<i>S. aureus</i>	21.3	62.5

Inhibition-zone values depend on the assay configuration and should be interpreted alongside appropriate controls.

In this example, *T. chebula* and *O. tenuiflorum* show the largest inhibition zones and the lowest MIC values. If a comparable pattern were observed in properly controlled experiments, it would justify further investigation of these extracts against additional microorganisms. However, a single-strain comparison cannot establish broad-spectrum antimicrobial activity.



Crude plant extracts may contain constituents with different diffusion rates and solubilities. Consequently, an extract with a relatively small inhibition zone may still contain compounds with measurable activity in broth dilution. Solvent controls are particularly important because residual extraction solvents can independently affect microbial growth.

Comparative anti-inflammatory activity

An in vitro protein-denaturation assay may provide preliminary information about the capacity of extracts to influence a model of protein destabilisation. Table 4 demonstrates a hypothetical comparison at a single extract concentration.

Table 4. Hypothetical inhibition of protein denaturation

Medicinal plant	Extract concentration (µg/mL)	Inhibition (%)
W. somnifera	200	52.4
T. cordifolia	200	61.8
O. tenuiflorum	200	68.5
A. racemosus	200	47.2
T. chebula	200	76.3

The illustrative results place T. chebula first and O. tenuiflorum second in percentage inhibition. Such an outcome would support further concentration–response testing, but it would not demonstrate that the extracts suppress inflammation through a specific molecular pathway. Direct measurement of inflammatory mediators, appropriate cytotoxicity testing and independent validation would be required before stronger mechanistic claims could be made.

DISCUSSION

Relationship between phytochemical composition and antioxidant activity

The proposed study integrates quantitative phytochemical measurements with antioxidant assays to investigate whether differences in chemical composition correspond to differences in radical-scavenging capacity. Phenolic compounds and flavonoids are plausible contributors because their chemical structures can support electron or hydrogen transfer reactions. Nevertheless, total phenolic content is an aggregate measure and does not identify individual compounds or establish their relative biological contribution.

The hypothetical dataset shows an inverse pattern between total phenolic content and DPPH IC₅₀. This relationship should be treated as a testable hypothesis rather than a finding. Folin–Ciocalteu measurements can also respond to non-phenolic reducing substances, while DPPH activity is affected by solvent, reaction time, steric accessibility and sample concentration. Complementary assays and chromatographic profiling would strengthen the interpretation.

Antimicrobial potential and experimental limitations

The proposed antimicrobial investigation addresses the possibility that plant-derived metabolites may inhibit microbial growth. Tannins, phenolics, terpenoids and

other constituents may interact with microbial structures or cellular processes, but the effects of crude extracts can be complex and synergistic or antagonistic interactions may occur.

The illustrative results indicate stronger activity for T. chebula and O. tenuiflorum against the selected bacterial strain. These patterns must not be generalised to other bacteria or fungi without direct testing. Moreover, MIC values from crude extracts do not provide the same clinical interpretation as susceptibility results for established antimicrobial drugs. Follow-up work should assess reproducibility, selectivity against microbial rather than mammalian cells, activity against additional reference strains and the stability of active fractions.

Anti-inflammatory potential and biological relevance

Protein-denaturation assays offer a relatively accessible screening approach but represent only a limited aspect of inflammatory biology. In living systems, inflammation involves interacting immune cells, signalling molecules, vascular responses and tissue-specific mechanisms. Inhibition of protein denaturation cannot establish modulation of these processes.

The hypothetical ranking of T. chebula and O. tenuiflorum provides an example of how candidate extracts might be selected for further study. A stronger evidence base would combine concentration-dependent results with cell-based inflammatory endpoints, measurements of cell viability and investigation of relevant pathways. Animal studies, where scientifically justified and ethically approved, would require separate protocols and should follow the principles of replacement, reduction and refinement.

Standardisation, reproducibility and safety

Plant identity, cultivation conditions, geographical origin, storage and extraction procedures can substantially influence chemical composition. Standardisation is therefore necessary both for experimental reproducibility and for any eventual development of a herbal product. Botanical authentication, voucher specimens, extract yields, physicochemical parameters and chromatographic fingerprints should be documented.

Safety must be investigated independently of efficacy. Traditional use does not guarantee that every preparation, dose or route of administration is safe. Contaminants, adulteration, interactions with medicines and intrinsic toxic constituents require consideration. The WHO guidance on medicinal-plant quality control provides a useful framework for establishing material identity and quality parameters.

LIMITATIONS OF THE STUDY

The proposed investigation has several methodological limitations. First, in vitro antioxidant assays do not directly predict antioxidant effects in humans because bioavailability, metabolism and tissue distribution



influence biological activity. Secondly, crude extracts contain mixtures of compounds, making it difficult to attribute observed responses to individual constituents. Thirdly, antimicrobial inhibition may be influenced by extract solubility and diffusion, while protein-denaturation assays provide only indirect evidence of anti-inflammatory potential.

The illustrative datasets presented in this paper are not substitutes for laboratory measurements. Consequently, no statistical significance, experimental reproducibility or biological superiority can be claimed from these values. Future work should employ authenticated samples from documented sources, validated protocols, appropriate controls, independent replication and transparent reporting of raw data. Chemical characterisation and toxicity testing should precede any consideration of therapeutic application.

CONCLUSION

The comparative evaluation of *Withania somnifera*, *Tinospora cordifolia*, *Ocimum tenuiflorum*, *Asparagus racemosus* and *Terminalia chebula* offers a structured approach to investigating the antioxidant, antimicrobial and anti-inflammatory potential of traditionally used medicinal plants. Integrating pharmacognostical authentication, phytochemical screening, quantitative chemical analysis and complementary biological assays can help identify promising extracts and establish a reproducible basis for further research.

The proposed framework also highlights the need to distinguish preliminary biological activity from demonstrated therapeutic efficacy. Antioxidant capacity measured through chemical assays, antimicrobial inhibition observed under laboratory conditions and responses in protein-denaturation models each address different experimental endpoints. Their combined interpretation requires appropriate controls, statistical analysis, chemical characterisation and safety assessment.

The hypothetical examples suggest how comparative tables and graphs may be structured, but they do not establish the actual performance of any selected plant. The principal contribution of this research framework is its emphasis on methodological consistency and evidence-based evaluation. Experimentally verified findings could contribute to the standardisation of medicinal plant materials and inform subsequent investigations into active constituents, mechanisms of action, safety and potential pharmaceutical applications.

REFERENCES

- [1]. Bag, A., Bhattacharyya, S. K., Pal, N. K., & Chattopadhyay, R. R. (2013). Anti-inflammatory, anti-lipid peroxidative, antioxidant and membrane stabilizing activities of hydroalcoholic extract of *Terminalia chebula* fruits. *Pharmaceutical Biology*, 51(12), 1515–1520.
- [2]. Khan, I., Ullah, Z., Shad, A. A., Fahim, M., & Öztürk, M. (2022). In vitro antioxidant, anticholinesterase inhibitory, and antimicrobial activity studies of *Terminalia chebula* (Retz) and *Terminalia arjuna* (Roxb). *South African Journal of Botany*, 146, 395–400.
- [3]. Chaiyana, W., et al. (2022). Anti-inflammatory and antimicrobial activities of fermented *Ocimum sanctum* Linn. extracts against skin and scalp microorganisms. *Chemistry & Biodiversity*.
- [4]. Chaudhary, A., Sharma, S., Mittal, A., Gupta, S., & others. (2020). Phytochemical and antioxidant profiling of *Ocimum sanctum*. *Journal of Food Science and Technology*, 57(10), 3852–3863.
- [5]. Saha, S., & Verma, R. J. (2016). Antioxidant activity of polyphenolic extract of *Terminalia chebula* Retz fruits. *Journal of Taibah University for Science*, 10(6), 805–812.
- [6]. Mandeville, A., & Cock, I. E. (2018). *Terminalia chebula* Retz. fruit extracts inhibit bacterial triggers of some autoimmune diseases and potentiate the activity of tetracycline. *Indian Journal of Microbiology*, 58(4), 496–506.
- [7]. In vitro assessment of antioxidant and antibacterial properties of *Terminalia chebula* Retz. leaves. *Asian Pacific Journal of Tropical Biomedicine*, 2(2, Suppl.), S788–S795.
- [8]. Gonfa, Y. H., Tessema, F. B., Bachheti, A., Rai, N., Tadesse, M. G., Singab, A. N., Chaubey, K. K., & Bachheti, R. K. (2023). Anti-inflammatory activity of phytochemicals from medicinal plants and their nanoparticles: A review. *Current Research in Biotechnology*, 6, Article 100152.
- [9]. Almatroodi, S. A., Alsahli, M. A., Almatroodi, A., & Rahmani, A. H. (2020). *Ocimum sanctum*: Role in diseases management through modulating various biological activity. *Pharmacognosy Journal*, 12(5), 1198–1205.